

Health Risk Factors

2022

JUNE 8

CARROLL COUNTY, KY



Introduction

Carroll County, KY

Health risk factors are the attributes, characteristics or exposures that increase the likelihood of a person developing a disease or health disorder. Risk factors are around us every day in our homes, places of work, schools, and in public. A nearby sneeze or cough may increase your likelihood of catching the flu, COVID-19, or a common cold. Being overweight increases the likelihood you'll develop diabetes and smoking increases your risks for many cancers.

To help understand the risk factors affecting the health of the populations in Carroll County we have collected and analyzed a wealth of health information for review and use.

“Carroll County, where rural and urban cultures meet”

This report will explore the health risk factors and pertinent details to help better understand how our populations health is being affected. With use of this information, it is anticipated, mitigation measures will be implemented to help improve the overall health of our communities.



Life Expectancy


Averages

Life expectancy is the average number of years a person is expected to live. The current U.S. life expectancy for persons born today is 79.05 years¹. Carroll County’s life expectancy is 72.51 years¹. While the overall rates have improved since 1980, the life expectancy in Carroll County remains considerably lower than the U.S. and state averages.

**72.5
years**

“The average resident dies 6.5 years sooner”

Some of the main factors affecting life expectancy include, but are not limited to, gender, genetics, prenatal and childhood conditions, education, socio-economic status, marital status, ethnicity/migrant status, lifestyle, and access to medical care/technology.



	Carroll, Kentucky	Lowest Life Expectancy in U.S. Ogala Lakota, SD	Highest Life Expectancy Summit, CO
Life Expectancy¹	72.5 years	64.5 years	91.72 years
Poverty Rate³	27.5%	46.8%	1.93%
Median Household Income³	\$43,524	\$31,423	\$80,709
Education Bachelor’s degree or higher 25+yrs⁴	7.4%	11.1%	53.6%

Carroll County comparison to the counties with the highest and lowest life expectancies in the U.S.


Life Expectancy

Gender


Like the state and nation overall, men have a lower life expectancy than women in Carroll County. Men’s overall life expectancy is 5.19 years less than women in Carroll County as compared to 5 and 5.09 in the Commonwealth and U.S. overall¹. There are several factors affecting the disparity between males and females. Some of the more obvious include the fact that men tend to take bigger risks than women resulting in higher accident rates. Additionally, men are also more likely to have jobs in riskier occupations such as military combat, firefighting, and working at construction sites where injury and accidents are more prevalent.

“On average, men die 5 years earlier than women”

Life Expectancy Trends - Males¹
Carroll County, Kentucky, U.S. 2000-2019




Life Expectancy Trends - Females¹
Carroll County, Kentucky, U.S. 2000-2019



Industry ⁵	Males	Females
Agriculture – forestry and mining	100%	0%
Construction	93.8%	6.2%
Utilities	87.3%	12.7%
Waste Mngt	95.4%	4.6%

Percentage of male to female workers by industry associated with more dangerous working conditions.

Life Expectancy Trends by Gender¹
Carroll County 2000-2019




Causes of Death

Leading Causes

Deaths are reported and tracked using a diagnosis code, currently the International Statistical Classification of Diseases and Related Health Problems, or (ICD10). These codes help in identifying trends and conducting analysis to identify causes of death in a population.

Since 1921, heart disease has been the leading cause of death in the U.S.⁶ In the years between 1900 and 1920 heart disease was second only to deaths caused by influenza/pneumonia and tuberculosis. The great influenza pandemic of 1918 that killed between 20 and 40 million people globally had a

**Top 10 Leading Causes of Death, Accumulative⁷
Carroll County 1999-2020**




material influence on those early rates. However, because of the introduction of antibiotics in the 1930s, the last 100 years has seen tuberculosis fall from a leading cause and influenza/pneumonia drop to the eighth leading cause of death in the U.S.⁷ Heart disease,

(for ICD10 codes I00-I09, I11, I13, I20-I51), however continues holding fast in its position as the leading cause of death for Americans⁷ resulting in more than 14 million deaths from 1999 to 2020⁷, or 4.3% of the total U.S. population today. As an accumulative total in the past 21 years, heart disease is considered a leading cause of death in Carroll County as well.

However, when examining the yearly trends in the leading causes of death, deaths from cancers overtook heart disease as the leading cause of death in 2009 in Carroll County and continues to hold that position as of 2020⁷.





**Leading Cause of Death Trend over Time⁷
Carroll County 1969-2020**



Causes of Death

Leading Causes by Age

Evaluating the leading cause of death by age group demonstrates, as expected, that the leading cause of death changes as we age. Persons between 15 and 44 years are more likely to die from primarily preventable causes such as motor vehicle accidents, accidental poisoning by narcotics and psychodysleptics, and suicide than any other causes.



*Deaths from motor vehicle accidents disproportionately affect those aged 15-44 in Carroll County by more than 4 times the national rate. #Lung Cancer kills Carroll County residents more than twice as frequently for those aged 45-64 and those aged 85+ die of Alzheimer's Disease 1.5 times more frequently than the average American.

Causes of Death

Underlying and Contributing Factors

When evaluating the cause of death, it is important to examine and understand the underlying and contributing factors that result in the diagnoses of specific diseases and causes of death. While genetics and environmental factors have a part, behavior plays a much larger role in the development of specific diseases that result in death than was believed in the past.

In 2009-2010, at least 47% of adults in the U.S. had at least one risk factor for heart disease⁸. According to the Centers for Disease Control and Prevention (CDC), there are three prevailing risk factors contributing to the incidence of heart disease including uncontrolled high blood pressure, uncontrolled high low-density lipoproteins cholesterol (LDL) and smoking. (The first two of which can be exacerbated by behavior and the last being solely a behavior.) Similarly, 90% of all lung cancers are the result of smoking tobacco⁹ and 80% of all patients diagnosed with COPD are or were smokers¹⁰.


An examination of the underlying causes of premature death (death before age 75)⁷, for Carroll County residents shows behavioral factors have a significant effect on the lower life expectancy trends as compared with the nation and state overall. Those behaviors including smoking, alcohol use, and drug/alcohol abuse¹¹.

ICD10	Description	Deaths per 100k ⁷	Underlying Contributing Factors/100k		
			Tobacco ¹¹	Alcohol ¹¹	Drug/Alcohol Abuse ¹¹
C34.9	Lung Cancer	71.4	64.3	2.5	
I21.9	Acute Mia	56.4	11.3		
J44.9	COPD	36.4	29.1		
I25.1	Atherosclerotic HD	22.7	4.5		
I50.0	Congestive HF	20.5	6.8		
V89.2	Motor Vehicle Accident	16.4		4.6	
C50.9	Breast Cancer	12.7		1.9	
X44	Accidental Poisoning	11.8			6.4
A41.9	Septicemia	9.6			
C18.9	Colon Cancer	9.1		.3	
J18.9	Pneumonia	9.1			
X70X74	Suicide	18.2			
X42	Narcotic Poisoning	9.1			4.9
E14.9	Diabetes w/o complications	8.2			
J44.0	COPD	7.7	6.2		
K70.3	Alcoholic Cirrhosis	7.7	7.7		
C80	Cancer unspecified	7.3			
J43.9	Emphysema	7.3	5.8		
I64	Stroke	5.9	1.2		
I11.9	Hypertensive HD	5.9	1.2		
C25.9	Cancer Pancreas	5.5		.2	
C92	Acute Myeloid Leukemia	5.0			
C64	Cancer Kidney	4.5	1.4		
I46.9	Cardiac Arrest, unspecified	4.5			

Cause of Death


Years of Potential Life Lost

Years of potential life lost, (YPLL), represent the years of life lost where death occurs before the age of 75, the average annual age of death for Americans. Carroll County has a very high YPLL that is increasing, as compared with state and top performers in the nation.



For those under the age of 75, the leading cause of death remains malignant neoplasms, 176.9 per 100k, followed closely by heart disease at 146.9 per 100k and accidents at 68.7 per 100k⁷.

Cause of Years of Potential Life Lost⁷
Carroll County 2022




Causes of Death

Cancers


Since the year 1999, cancer of the lungs or bronchus has represented the leading cause of death among persons diagnosed with malignancies in Carroll County, accounting for 95.1 deaths per 100,000 population⁷.

Leading Cancer Cause of Death⁷
Carroll County 1999-2020



Lung cancer is the cause of more than half of all cancer related deaths in Carroll County⁷. According to the CDC, smoking can cause cancers almost anywhere in the body including the bladder, blood, cervix, colon, rectum, esophagus, kidney, renal pelvis, liver, lungs, bronchi, trachea, mouth, throat, pancreas, stomach and larynx¹³. Additionally, Lung Cancer has a known association with the inhalation of radon gas. Higher cancer rates in Carroll County may be the result of either or both underlying contributing factors. See page 19 for details on tobacco rates. See page 24 for details on radon in Carroll County.

Leading Cancer Cause Death⁷
Three Rivers District 1999-2020



ICD10	Cancer
C34.9	Lung
C18.9	Colon
C50.9	Breast
C80	Unspecified
C61	Prostate
C25.9	Pancreas
C64	Kidney
C67.9	Bladder
C56	Ovarian
C15.9	Esophagus
C85.9	Lymphoma
C71.9	Brain
C90.0	Multiple Myeloma




Health Conditions

Asthma

According to the CDC, 1 in 13 persons in the U.S. has asthma¹⁴. Most people can control their symptoms and prevent asthma attacks by avoiding triggers and using prescribed medication such as inhalants correctly.

Asthma Prevalence^{14,15}

Carroll County, Kentucky, U.S. 2016-2018





Carroll County has a higher-than-average prevalence for asthma at 12.3%, or 1 in 8, when compared with the state and national averages at 11.5% and 7.7%¹⁵.

“1 in 8 people in Carroll County has Asthma”

Asthma induced ER visits in Carroll County in 2019 represent an equivalent of 3.4% of the total population or 28% of those with Asthma¹⁶.

Asthma induced Emergency Room Visits Age Adjusted Fine Particulate Air Matter (Modeled)¹⁶
Carroll County & Kentucky 2008-2019






Health Conditions

Chronic Obstructive Pulmonary Disease

Chronic Obstructive Pulmonary Disease, or COPD, is a group of pulmonary disease that restricts airflow within the lungs and includes chronic bronchitis and emphysema. According to the CDC, approximately 18 million Americans have been diagnosed with


COPD induced Hospitalizations Age Adjusted¹⁶
Carroll County & Kentucky 2008-2019



COPD and millions more suffer with it but have yet to be diagnosed and receive treatment¹⁷. People aged 65 and over are more likely to be diagnosed with some form of COPD but it can affect persons of any age.

Nationally, there are 40.4 ER visits per 100 persons and 43.5% of those visits are less than 15 minutes in duration¹⁸. According to the National Hospital Ambulatory Medical Care Survey: 2018 Emergency Department Summary Tables, COPD accounted for 8.1 million ER visits in 2018 or 6.2% of all ER visits nationwide¹⁸. Carroll County has a very high rate of ER visits as compared with the state which may have multiple causes including the lack of access to preventive and acute medical care.

COPD induced Emergency Room Visits Age Adjusted¹⁶
Carroll County & Kentucky 2008-2019




Health Conditions

Heart Attacks




A heart attack or myocardial infarction is a result of reduction or block of blood flow to the heart that results in the damage or death of part of the heart muscle. Coronary Artery Disease (CAD) is the primary cause of myocardial infarction. Coronary Artery Disease is caused by the buildup of plaque and other deposits in the artery(ies) creating narrowing, reducing blood flow and supply to critical body parts/organs.

**Heart Attack induced Hospitalizations Age Adjusted¹⁹
Carroll County & Kentucky 2008-2019**



“Carroll County has 1.6 times more heart attacks than KY.”

When overlaying mortality associated with heart attacks in Carroll County, the



percentages of heart attacks resulting in death is high⁷.

(Note, years with deaths below a specific threshold are suppressed to protect privacy and are not displayed.)


Health Conditions

Diabetes Prevalence

According to the CDC, 37.3 million people have diabetes in the U.S. and 23% are undiagnosed²⁰. Diabetes is a chronic life-long disease characterized by high blood sugar resulting in the 7th leading cause of death in the U.S. It is the number one cause of kidney failure, lower-limb amputations and adult blindness²¹.

Type 2 diabetes is the most common form of the disease affecting between 90-95% of those diagnosed with diabetes²². It may be prevented or delayed with healthy lifestyle changes such as losing weight, eating healthy foods, and being active.

As of 2019, 1 in 13 people in Carroll County have been diagnosed with diabetes and males have slightly higher rates of diabetes diagnoses than females.




Risks for Pre-Diabetes and type 2 diabetes²³

- Overweight
- 45 and older
- Parent or sibling with type 2 diabetes
- Physically active less than 3x/week
- Had gestational diabetes
- African American, Hispanic/Latino American, Pacific Islander, American Indian, or Asian American


Health Conditions

Sexually Transmitted Infections

Carroll County has maintained a similar rate of Chlamydia as the state overall. Periodically the incidence peaks above the state rate but consistently remains lower with an eight-year trend of 376.1 per 100k persons²⁵ compared to 419.4 in the state overall. Gonorrhea incidence remains significantly lower than the state rate except for the year 2012 where the rate was nearly the same as the state and in 2018 where it exceeded the state rate by 10 persons per 100k²⁵.



“47.4% of youths did not use a condom in their last encounter”



Approximately 40.4% of Kentucky high school students have had at least one sexual encounter²⁶. Of those, 7.2% had sexual intercourse with four or more persons and 30.7% are currently sexually active²⁶.


Health Conditions

Overweight and Obesity – Adults



Adult obesity is defined as a weight that is higher than what is considered healthy for a person’s height. In the years 2021, the adult obesity prevalence was 41.9% of U.S. adults.²⁷

Kentucky leads the nation in obesity rates with obesity rates averaging 40-45% of the population overall.




“63% of Ky counties have lower obesity rates than Carroll Co.”

The most recent county level data in Carroll County reflect obesity rates that align with state prevalence rates at 39.9% of the total population in 2020²⁸.

Obesity age-adjusted prevalence (%)


- > 42.8 - 46.2
- > 40.6 - 42.8
- > 38.8 - 40.6
- > 36.6 - 38.8
- 33.2 - 36.6



Health Conditions


Overweight and Obesity by Race and Ethnicity

Obesity prevalence affects persons of varying races and ethnicities differently among populations. The prevalence of Self-Reported Obesity, 2019-2021 Behavioral Risk Factor Surveillance System (BRFSS) reflects significant differences in obesity prevalence rates among non-Hispanic, Hispanic, black, white, and Asian populations. Non-Hispanic Asian adults have the lowest obesity prevalence at just 13.8% of the U.S. population as compared with all other races and Hispanic populations²⁹.



In Carroll County, 1 in 3.2 White Adults; 1 in 2.8 Hispanic adults; 1 in 2.5 Black adults and 1 in 7.2 Asian adults are obese²⁹.

Age-adjusted, nationally applied rate of prevalence of adult obesity by race/ethnicity, based on Ky Averages²⁹
Carroll County, 2020




Health Conditions

Childhood Obesity

Childhood overweight is defined as a Body Mass Index (BMI) at or above the 85th percentile and below the 95th percentile for children and teens of the same age and sex. Childhood obesity is defined as a BMI at or above the 95th percentile for children and teens of the same age and sex. The National Health and Nutrition Examination Survey 2017-March 2020 Pre-pandemic, indicates an overall obesity prevalence rate among youth aged 2-19 years of 19.7% in the U.S.³⁰ The survey also found that obesity rates increased among youth as age increased.


Youth Obesity Rate by Age Group³⁰
U.S. 2017-March 2020



“Recent studies show childhood obesity rates are increasing”

As an attempt to help slow the spread of COVID-19 in Kentucky schools, the last two school years, 2020-2021 and 2021-2022 did not conduct mass health screenings in Kentucky. Therefore, local BMI data for Carroll County schools is extremely limited and has a very low confidence interval and has not been included in this most recent report. It is worth noting however that the average percent of children considered

Youth Obesity Rate Trends³¹
Kentucky Ages 10-17, 2016-2020



obese in Kentucky schools for ages 10-17 has increased from 19.3% of the population in the school year 2016-2017 to 23.8% in the school year 2019-2020³¹.




Health Risk Behaviors

Diet

Healthy behaviors such as a healthy diet and regular physical activity contribute to better short term health outcomes, long term health benefits, and the prevention of chronic diseases such as Type 2 diabetes and heart disease.

Factors affecting healthy behaviors related to diet and exercise include access to and availability of healthy foods and opportunities for exercise. Access is limited in populations that are low income, have limited grocery supermarkets, have a high fast-food concentration and lack of access to transportation. The U.S. Department of Agriculture Food Access Research Atlas indicators reveal a significant portion of Carroll County is categorized as low income³². Additionally, 5.3% of home in the census tract 21041950200, which includes Carrollton, has low vehicle access and are more than 1/2 a mile from a supermarket³².



Carroll County is at a significant disadvantage when compared with residents in areas such as Boone, Kenton, Campbell, and Oldham counties that have little to no low-income areas and limited populations without access to transportation.

Food establishments³³
Groceries – 3
Full-Service Restaurant – 5
Fast Food Restaurant – 13
Convenience Stores – 5
Famer’s Market - 1


Health Risk Behaviors

Exercise

Physical inactivity is a critical factor in the health status of a population. Exercise has been linked to healthy weight and BMI, reduction in various chronic diseases, normal growth and development, better quality of life, and healthy sleep patterns.

“More than 35% of Carroll County residents are inactive”

The CDC U.S. Diabetes Atlas³⁴ reports Carroll County has a self-reported “lack of physical activity” that is statistically higher than the state overall indicating they are less physically active than the average Kentuckian. As of 2020, Carroll County has a physical inactivity rate of 35.8% of the population reporting no leisure time physical activity in the last 30 days³⁴.





The source of data used in the Atlas includes the BRFSS data which defines physical inactivity as those respondents who answered no to the question “During the past month, other than your job, did you participate in any physical activities or exercises or exercises such as running, calisthenics, golf, gardening, or walking for exercise?”

Health Risk Behaviors

Tobacco Use Adults

Smoking remains the single largest leading cause of preventable disease, death, and disability, in the U.S., accounting for more than 480,000 deaths every year, or about 1 in 5 deaths³⁵. The Centers for Disease Control and Prevention reports that adult smoking rates have declined in the U.S. from 1 out of nearly every 5³⁵ adults in 2005 to 1 in 10 as of 2020³⁶.

Carroll County has a long history of tobacco as a means of living for many families. This history has included husband and wife working side by side in the fields along with their children for many decades and the annual “Tobacco Festival” which is entering its 73rd year in 2022. Expectedly, Carroll County has a significantly higher rate of adults who smoke, with a rate of 28% or 1 in 4 adults³⁷.



The financial burden of tobacco use is significant, costing the U.S. billions of dollars each year. Carroll County’s long history with tobacco as an economic driver and familial heritage, presents an especially difficult challenge as it pertains to health status and public opinion of tobacco use and comprehensive tobacco free policies.

Health Risk Behaviors



Tobacco Use Adolescents

The Centers for Disease Control and Prevention reports that 1 in 4 High School Students used some form of tobacco product in 2022 with the greatest percentage of those using e-cigarettes in the past 30 days at 14.1%³⁸.

“As many as 1 in 6 HS students report using Tobacco”

Today, girls are more likely to report using of some tobacco product in the last 30 days than boys, a shift since 2018 when more boys reported use than girls, with E-Cigarettes representing 15.4% of tobacco use in girls³⁸. The same survey found similar findings in middle school students reporting use of some form of tobacco product with most of these using e-cigarettes (4.1%)³⁸.

Assuming similar applied ratios of tobacco use in the youth population as compared with national averages, middle and high school use could be as high as 1 in 12 or as low as 1 in 22 for middle school students and as high as 1 in 3 or as low as 1 in 6 for high school students in Carroll County.




Health Risk Behaviors

Alcohol Use

In the years 2015-2019 excessive alcohol use was responsible for more than 140,000 deaths in the U.S. each year, or more than 380 deaths per day³⁹. In Kentucky, men with alcohol related deaths were nearly four times more likely to have chronic alcohol abuse than women⁴⁰.

According to the BRFSS survey results as reported in the 2022 County Health Rankings, Carroll County has a slightly lower than average number of adults reporting binge or heavy drinking in the last 30 days that the state overall⁴¹. It should be noted that data prior to 2016 is not available for Carroll County.



Excessive or heavy drinking is associated with short and long term health effects including: injuries such as motor vehicle crashes, falls, drownings, and burns; violence including homicide, suicide, sexual assault, and intimate partner violence; alcohol poisoning, risky sexual behaviors including unprotected sex with multiple partners which can result in unwanted pregnancies or sexually transmitted diseases including HIV; and miscarriage or still birth, or fetal alcohol spectrum disorders among pregnant women.

Long term health risks include high blood pressure; heart disease, stroke, liver disease, and digestive problems; cancer of the breast, mouth, throat, esophagus, liver, and colon; learning and memory problems including dementia; mental health problems including depression and anxiety; social problems including lost productivity, family problems and unemployment; alcohol dependence or alcoholism.

Excessive alcohol use includes binge drinking, heavy drinking, any alcohol use by persons under 21 years and alcohol used by pregnant women.

Binge drinking is defined as 5 or more drinks consumed on a single occasion for men and 4 or more for women, generally within about 2 hours.


Heavy drinking is defined as men who consume 15 drinks or more per week and women who consume 8 drinks or more per week.

Health Risk Behaviors

Recreational Drug Use/Abuse

Carroll County has been ranked 67th for the most vulnerable counties in the nation at risk for an outbreak of HIV and/or Hepatitis C (HEP C)⁴² among persons who inject drugs. Of the 120 counties in Kentucky, 36 are ranked in the top 100 at risk, and 54, or nearly half, are in the top 220 counties across the nation identified at risk⁴². One of the primary objectives of the [Hep C Elimination Plan in Kentucky](#) is to increase and improve equitable access to harm reduction services across social and demographic areas particularly among people who use drugs. Along the I-71 corridor from Northern Kentucky to


Jefferson County, there are no Syringe programs until you arrive in the Louisville area⁴³.



According to the 2021 Kentucky Drug Overdose Deaths Report, Carroll County had an age adjusted mortality rate of 111.88 per 100,000 persons⁴⁴.

2021 Drug Overdose Deaths Kentucky⁴⁴


- Suppressed
- 0 per 100k
- 100 - 110 per 100k
- 111 - 125 per 100k
- 126 - 150 per 100k
- Less than 100 per 100k




HEALTH RISK FACTORS

Overdose Mortality

In Carroll County, deaths associated with drug overdose rates decreased 13.34% from 62.2 to 53.9 per 100k, between the 2011-2015 and 2016-2020 time periods⁴⁵. Compared with the Commonwealth and the U.S. Carroll County mortality from drug overdoses is slightly lower than the Commonwealth but remains higher than the national rate at 53.9 compared with 32.5⁴⁵.



Deaths from opioid specific overdoses also decreased but only slightly, representing a 1.24% decrease between the 2011-2015 and 2016-2020 time periods⁴⁵. In contrast, rates for opioid specific drug overdose deaths in the Commonwealth and U.S. both increased during the same time period that Carroll County experienced a decrease⁴⁵.




Environmental Factors

Air Particulate Matter

Per the CDC, Environmental Public Health Tracking Network, air pollution poses a public health threat affecting potentially millions of people throughout the U.S.⁴⁶ Pollution such as dust, dirt, soot, and smoke are one kind of air pollution called fine particulate matter (FPM). Sources of FPM include forest fires, power plants, industries, and automobiles. FPM is associated with compromised health and may lead to breathing problems, worsening of asthma symptoms and/or worsening of some heart conditions, as well as increased low birth weight⁴⁷.

Decreased lung function, chronic bronchitis, asthma, and adverse pulmonary effects are common in populations exposed to high levels of FPM. Long term exposure is associated with premature death of people aged 65 and older even when exposure is at levels below the National Standards⁴⁸. Since 2001 FPM rates in Carroll County have

decreased by 42% to 9.1 PM ug/M³ in 2018⁴⁹.




Rates below 12 Ug/m³ are considered acceptable according to the U.S. Environmental Protection Agency

Environmental Factors

Radon

Radon is a gas formed naturally in the environment with the breakdown of radioactive metals. People are always exposed to radon because it comes from the earth, but some areas have higher concentrations of the gas than others. The gas can enter your home through cracks and gaps.

Kentucky has an average radon level of 5.4 picocuries per liter of air (pCi/L)⁵⁰. The




Environmental Protection Agency indicates remediation should be taken for levels exceeding 4pCi/L in a home⁵¹. Higher radon levels are associated with increased risk of lung cancer and is the second leading cause of all lung cancers in the U.S.

“The average Carroll County radon level for homes tested is 2.6pCi/L”

Of the homes tested, the average indoor radon level is 2.6 pCi/L as compared to 1.3pCi/L nationally⁵². Carroll County homes have higher indoor radon levels than the

**Indoor Radon Levels measured in pCi/L⁵²
Carroll County, 2022**



nation. Of the homes measured in Carroll County, 19% have radon levels of 4pCi/L or higher. More than half of the homes in Carroll County have insufficient data to measure levels⁵².

End Notes

References and Data Sources

1. Institute of Health Metrics Evaluation, US Health Map, Viz Hub, Life Expectancy, 2000-2019 as accessed at <https://vizhub.healthdata.org/subnational/usa>
2. Institute of Health Metrics Evaluation, 1980-2014 United States Life Expectancy and Age-specific Mortality Risk by County 1980-2014. Seattle, United States of America, institutes for Health Metrics and Evaluation (IHME), 2017, as accessed at <https://ghdx.healthdata.org/record/ihme-data/united-states-life-expectancy-and-age-specific-mortality-risk-county-1980-2014>
3. U.S. Census Bureau, American Fact Finder, 2020 ACS 5 Year Estimates, Table DP03 – Selected Economic Characteristics
4. U.S. Census Bureau, American Fact finder, 2020 ACS 5 Year Estimates, Table S1501 – Educational Attainment
5. U.S. Census Bureau, American Fact Finder, 2020 ACS 5 Year Estimates, Table S2404 – Industry by Sex for the Full-Time, Year-Round Civilian Employed Population 16 Years and Over
6. Centers for Disease Control and Prevention, National Center for Health Statistics, Leading Causes of Death, 1900-1998 as accessed at https://www.cdc.gov/nchs/data/dvs/lead1900_98.pdf
7. Centers for Disease Control and Prevention, National Center for Health Statistics. National Vital Statistics System, Mortality 1999-2020 on CDC WONDER Online Database, released in 2021. Data are from the Multiple Cause of Death Files, 1999-2020, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program. Accessed at <http://wonder.cdc.gov/ucd-icd10.html> on Dec 5, 2022, 2:24:37 PM
8. Fryar CD, Chen T, Li X. Prevalence of uncontrolled risk factors for cardiovascular disease: United States, 1999–2010. NCHS data brief, no 103. Hyattsville, MD: National Center for Health Statistics. 2012. As accessed at <https://www.cdc.gov/nchs/data/databriefs/db103.htm>
9. U.S. Department of Health and Human Services. The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General. Atlanta: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2014 As accessed at: https://www.cdc.gov/tobacco/data_statistics/sgr/50th-anniversary/index.htm#report
10. The COPD Foundation, Understanding COPD, What Causes COPD. As accessed at <https://www.copdfoundation.org/What-is-COPD/Understanding-COPD/What-Causes-COPD.aspx>
11. See Appendix A
12. County Health Rankings, Carroll County, Kentucky, Premature Death, 2022.
13. The Centers for Disease Control and Prevention, Smoking and Cancer, May 2022, as accessed at: <https://www.cdc.gov/tobacco/campaign/tips/diseases/cancer.html>
14. The Centers for Disease Control and Prevention, National Current Asthma Prevalence, 2020, as accessed at: https://www.cdc.gov/asthma/most_recent_national_asthma_data.htm
15. The Centers for Disease Control and Prevention, State Maps for Asthma Prevalence by Six-Level Urban-Rural Classification, 2016-2018, as accessed at: <https://www.cdc.gov/asthma/national-surveillance-data/asthma-prevalence-state-classification.htm>
16. Kentucky Environmental Health Tracking, Asthma induced Emergency Room Visits Age Adjusted and Fine Particulate Air Matter Modeled as access at: <https://kyibis.mc.uky.edu/ehl/>
17. Centers for Disease Control and Prevention, National Trends COPD, As accessed at: <https://www.cdc.gov/copd/data-and-statistics/national-trends.html>
18. National Hospital Ambulatory Medical Care Survey: 2018 Emergency Department Summary Tables as accessed at: https://www.cdc.gov/nchs/data/nhamcs/web_tables/2018-ed-web-tables-508.pdf

19. Kentucky Environmental Health Tracking, Number of Hospitalizations for Heart Attack by County, as access at: <https://kyibis.mc.uky.edu/ehl/>
20. Centers for Disease Control and Prevention, Fast Facts on Diabetes as accessed at: <https://www.cdc.gov/diabetes/data/statistics-report/index.html>
21. Centers for Disease Control and Prevention, Diabetes Basics, What is Diabetes, as accessed at: <https://www.cdc.gov/diabetes/basics/diabetes.html>
22. Center for Disease Control and Prevention, Diabetes Basics, Type 2 Diabetes, as accessed at: <https://www.cdc.gov/diabetes/basics/type2.html>
23. Centers for Disease Control and Prevention, About Prediabetes & Type 2 Diabetes as accessed at: <https://www.cdc.gov/diabetes/prevention/about-prediabetes.html>
24. Centers for Disease Control and Prevention, United States Diabetes Surveillance System, Surveillance, Diabetes Prevalence, Carroll County as accessed at: <https://gis.cdc.gov/grasp/diabetes/DiabetesAtlas.html>
25. Centers for Disease Control and Prevention, Atlas Plus, HIV, Hepatitis, STD, TB, SDOH, 2020 as accessed at: <https://gis.cdc.gov/grasp/nchhstpatlas/tables.html>
26. Youth Behavior Risk Surveillance Survey 2019, Kentucky Results, as accessed at: <https://nccd.cdc.gov/youthonline/App/Results.aspx?LID=KY>
27. Centers for Disease Control and Prevention, Adult Obesity, as accessed at: <https://www.cdc.gov/obesity/data/adult.html>
28. Centers for Disease Control and Prevention, Explore Places Interactive Map, Obesity Rates 2020, as accessed at: https://experience.arcgis.com/experience/dc15b033b88e423d85808ce04bd7a497/page/Health-Outcomes/?data_id=dataSource_17-PLACES_LocalData_for_BetterHealth_5583%3A1140&views=Obesity
29. Centers for Disease Control and Prevention, Overweight & Obesity, Data & Statistics, Adult Obesity Prevalence Maps as accessed at: <https://www.cdc.gov/obesity/data/prevalence-maps.html#race>
30. National Health and Nutrition Examination Survey 2017-March 2020 Pre-pandemic Data Files Development of Files and Prevalence Estimates for Selected Health Outcomes, Youth Obesity, As accessed at: <https://stacks.cdc.gov/view/cdc/106273>
31. Robert Wood Johnson Foundation, State of Childhood Obesity, Explore State Data, Kentucky, Children ages 10-17, as accessed at: <https://stateofchildhoodobesity.org/state-data/?state=ky>
32. U.S. Economic Research Service, U.S. Department of Agriculture, Food Access Research Atlas as accessed at: <https://www.ers.usda.gov/data-products/food-access-research-atlas/go-to-the-atlas/>
33. U.S. Department of Agriculture, Food Environment Atlas, Carroll County. As accessed at: <https://www.ers.usda.gov/data-products/food-environment-atlas/go-to-the-atlas/>
34. Centers for Disease Control and Prevention, Explore Places Interactive Map, Physical Inactivity Rates 2020, as accessed at: https://experience.arcgis.com/experience/dc15b033b88e423d85808ce04bd7a497/page/Health-Outcomes/?data_id=dataSource_17-PLACES_LocalData_for_BetterHealth_5583%3A1140&views=Obesity
35. U.S. Department of Health and Human Services. The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General. Atlanta: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2014. As accessed at: https://www.ncbi.nlm.nih.gov/books/NBK179276/pdf/Bookshelf_NBK179276.pdf
36. Centers for Disease Control and Prevention, Smoking & Tobacco Use, Data and Statistics, Fast Facts and Fact Sheets, Smoking and Cigarettes as accessed at: https://www.cdc.gov/tobacco/data_statistics/fact_sheets/fast_facts/index.htm?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Ftobacco%2Fdata_statistics%2Ffact_sheets%2Findex.htm
37. County Health Rankings, Kentucky, Adult Smoking, 2022 as accessed at: <https://www.countyhealthrankings.org/explore-health-rankings/kentucky/carroll?year=2020>

38. Centers for Disease Control and Prevention, Tobacco, Data and Statistics, Fact Sheets, Youth and Tobacco Use, as accessed at: https://www.cdc.gov/tobacco/data_statistics/fact_sheets/youth_data/tobacco_use/index.htm
39. Centers for Disease Control and Prevention, Alcohol and Public Health, Data on Excessive Drinking as accessed at: <https://www.cdc.gov/alcohol/data-stats.htm>
40. Centers for Disease Control and Prevention, Alcohol-Related Disease Impact (ARDI), Alcohol-attributable deaths, 2015-2019 as accessed at: https://nccd.cdc.gov/DPH_ARDI/Default/Default.aspx
41. County Health Rankings, Carroll County, Kentucky, Excessive Drinking, 2022 as accessed at: <https://www.countyhealthrankings.org/explore-health-rankings/kentucky/carroll?year=2022>
42. Centers for Disease Control and Prevention, Vulnerable Counties and Jurisdictions Experiencing or At-Risk of Outbreaks, as accessed at: <https://www.cdc.gov/pwid/vulnerable-counties-data.html>
43. Kentucky Cabinet for Health and Family Services, Department for Public Health, Division of Epidemiology and health Planning, HIV/AIDS Branch, Syringe Exchange Programs as accessed at: <https://www.chfs.ky.gov/agencies/dph/dehp/hab/Pages/kyseps.aspx>
44. Kentucky Office of Drug Control Policy, 2021 Overdose Fatality Report as accessed at: [https://odcp.ky.gov/Reports/2021%20Overdose%20Fatality%20Report%20\(final\).pdf](https://odcp.ky.gov/Reports/2021%20Overdose%20Fatality%20Report%20(final).pdf)
45. Recovery Ecosystem Index as accessed at: https://rsconnect.norc.org/recovery_ecosystem_index/
46. Centers for Disease Control and Prevention, National Environmental Public Health Tracking, Explore Environmental Health Data, Air Quality as accessed at: <https://www.cdc.gov/nceh/tracking/topics/AirQuality.htm>
47. Particle Pollution, Fine particulate Air Matter, Centers for Disease Control and Prevention. As accessed at: https://www.cdc.gov/air/particulate_matter.html
48. The Harvard T.H. Chan School of Public Health, Harvard Data Science Initiative, Press Release, 2017. As accessed at: <https://www.hsph.harvard.edu/news/press-releases/u-s-seniors-air-pollution-premature-death/>
49. Centers for Disease Control and Prevention National Environmental Public Health Tracking Network, Fine Particulate Air Matter as accessed at: <https://ephtracking.cdc.gov/DataExplorer/>
50. American Lung Association, Radon Testing Disparities, Kentucky, Radon Testing in the State of Kentucky as accessed at: <https://www.lung.org/getmedia/5e0f8076-92c9-482e-b658-588d97b0b7e5/Radon-Testing-Disparity-report-for-Kentucky-2022-08-14>
51. U.S. Environmental Protection Agency, A Citizen's Guide to Radon, Protecting yourself and your Family from Radon, as accessed at: https://www.epa.gov/sites/default/files/2016-12/documents/2016_a_citizens_guide_to_radon.pdf
52. University of Kentucky, Kentucky Geological Survey, Radon as accessed at: <https://www.uky.edu/KGS/radon/>

Appendix A

ICD10	Tobacco Association	Alcohol Association
C34.9	https://pubmed.ncbi.nlm.nih.gov/16100660/	https://www.livescience.com/27259-alcohol-causes-cancer-deaths.html
I21.9	https://www.ncbi.nlm.nih.gov/books/NBK53012/	
J44.9	https://www.cdc.gov/tobacco/data_statistics/fact_sheets/health_effects/effects_cig_smoking/index.htm	
I25.1	https://www.ncbi.nlm.nih.gov/books/NBK53012/	
I50.0	https://www.ncbi.nlm.nih.gov/books/NBK53012/	
V89.2		https://www.cdc.gov/motorvehiclesafety/impaired_driving/impaired_drv_factsheet.html#statistics
C50.9		https://www.livescience.com/27259-alcohol-causes-cancer-deaths.html
X44		CDC wonder calculated by # of deaths caused by drug and alcohol induced from total poisoning deaths
A41.9		
C18.9		https://www.livescience.com/27259-alcohol-causes-cancer-deaths.html
J18.9		
X70/X74		
X42		CDC wonder calculated by # of deaths caused by drug and alcohol from total poisoning deaths
E14.9		
J44.0	https://www.cdc.gov/tobacco/data_statistics/fact_sheets/health_effects/effects_cig_smoking/index.htm	
K70.3		
C80		
J43.9	https://www.cdc.gov/tobacco/data_statistics/fact_sheets/health_effects/effects_cig_smoking/index.htm	
I64	https://www.ncbi.nlm.nih.gov/books/NBK53012/	
I11.9	https://www.ncbi.nlm.nih.gov/books/NBK53012/	
C25.9		https://www.livescience.com/27259-alcohol-causes-cancer-deaths.html
C92		
C64	https://www.cancer.net/cancer-types/kidney-cancer/risk-factors-and-prevention#:~:text=Smoking.,Gender.	
I46.9		