

# Health Risk Factors

2022

JUNE 8

PENDLETON COUNTY, KY



# Introduction

Pendleton County, KY

Health risk factors are the attributes, characteristics or exposures that increase the likelihood of a person developing a disease or health disorder. Risk factors are around us every day in our homes, places of work, schools, and in public. A nearby sneeze or cough may increase your likelihood of catching the flu, COVID-19, or a common cold. Being overweight increases the likelihood you'll develop diabetes and smoking increases your risks for many cancers.

To help understand the risk factors affecting the health of the populations in Pendleton County, we have collected and analyzed a wealth of health information for review and use.

***“Pendleton County, where community is about the people”***

This report will explore the health risk factors and pertinent details to help better understand how our populations health is being affected. With use of this information, it is anticipated, mitigation measures will be implemented to help improve the overall health of our communities.



# Life Expectancy


## Averages

Life expectancy is the average number of years a person is expected to live. The current U.S. life expectancy for persons born today is 79.05 years<sup>1</sup>. Pendleton County's life expectancy is 74.35 years<sup>1</sup>. Overall rates have increased slightly since 1980, increasing 2.15 years or 2.98%. When compared with the state and national life expectancy increases Pendleton County's life expectancy has increased at a much slower rate with state and national increase rates at 4% and 7.11%<sup>1</sup>. Accordingly, the life expectancy in Pendleton County remains considerably lower than the US and state averages.

**74.35  
years**

*“The average resident dies 4.7 years sooner”*

Some of the main factors affecting life expectancy include but are not limited to gender, genetics, prenatal and childhood conditions, education, socio-economic status, marital status, ethnicity/migrant status, lifestyle, and access to medical care/technology.



	Pendleton, Kentucky	Lowest Life Expectancy in U.S. Ogala Lakota, SD	Highest Life Expectancy Summit, CO
<b>Life Expectancy<sup>1</sup></b>	74.35 years	64.5 years	91.72 years
<b>Poverty Rate<sup>3</sup></b>	10.9%	46.8%	1.93%
<b>Median Household Income<sup>3</sup></b>	\$56,000	\$31,423	\$80,709
<b>Education Bachelor's degree or higher 25+yrs<sup>4</sup></b>	14.2%	11.1%	53.6%

*Pendleton County comparison to the counties with the highest and lowest life expectancies in the U.S.*

# Life Expectancy


## Gender




Like the state and nation overall, men have a lower life expectancy than women in Pendleton County. Men’s overall life expectancy is 5.19 years less than women in Pendleton County, as compared to 5 and 5.09 in the Commonwealth and U.S. overall<sup>1</sup>. There are several factors affecting the disparity between males and females. Some of the more obvious include the fact that men tend to take bigger risks than women resulting in higher accident rates. Additionally, men are also more likely to have jobs in riskier occupations such as military combat, firefighting, and working at construction sites where injury and accidents are more prevalent.

*“On average, men die 5+ years earlier than women”*

**Life Expectancy Trends - Males<sup>1</sup>**  
Pendleton County, Kentucky, U.S. 2000-2019




**Life Expectancy Trends - Females<sup>1</sup>**  
Pendleton County, Kentucky, U.S. 2000-2019



Industry <sup>5</sup>	Males	Females
Agriculture/Forestry	0%	100%
Mining	100%	0%
Construction	100%	0%
Utilities	18%	82%
Waste Mngt	53.9%	46.1%

Percentage of male to female workers by industry associated with more dangerous working conditions.

**Life Expectancy Trends by Gender<sup>1</sup>**  
Pendleton County 2000-2019




# Causes of Death

## Leading Causes

Deaths are reported and tracked using a diagnosis code, currently the International Statistical Classification of Diseases and Related Health Problems, or (ICD10). These codes help in identifying trends and conducting analysis to identify causes of death in a population.

Since 1921, heart disease has been the leading cause of death in the U.S.<sup>6</sup> In the years between 1900 and 1920 heart disease was second only to deaths caused by influenza/pneumonia and tuberculosis. The great influenza pandemic of 1918 that killed between 20 and 40 million people globally had a

**Top 10 Leading Causes of Death, Accumulative<sup>7</sup>  
Pendleton County 1999-2020**




material influence on those early rates. However, because of the introduction of antibiotics in the 1930s, the last 100 years has seen tuberculosis fall from a leading cause and influenza/pneumonia drop to the eighth leading cause of death in the U.S.<sup>7</sup> Heart disease,

(for ICD10 codes I00-I09, I11, I13, I20-I51), however continues holding fast in its position as the leading cause of death for Americans<sup>7</sup> resulting in more than 14 million deaths from 1999 to 2020<sup>7</sup>, or 4.3% of the total U.S. population today. As an accumulative total in the past 21 years, heart disease is considered a leading cause of death in Pendleton County as well.

However, when examining the yearly trends in the leading causes of death, the gap between deaths from cancers and heart disease has narrowed to the point it is only 1/10<sup>th</sup> of a decimal point difference<sup>7</sup>. Like neighboring communities, deaths from cancer are set to become the leading cause of death in Pendleton County. It is worth noting that the cancer related deaths have remained relatively steady since 1969.





**Leading Cause of Death Trend over Time<sup>7</sup>  
Pendleton County 1969-2020**



# Causes of Death

## Leading Causes by Age

Evaluating the leading cause of death by age group demonstrates, as expected, that the leading cause of death changes as we age. Persons between 15 and 44 years are more likely to die from primarily preventable causes such as motor vehicle accidents, accidental poisoning by narcotics and psychodysleptics, and suicide than any other causes.



\*Deaths from motor vehicle accidents disproportionately affect those aged 15-44 in Pendleton County by more than 5.6 times the national rate. #Congestive Heart Failure kills Pendleton County residents more than five times as frequently for those aged 45-64 and those aged 85+ die of Cardiac Arrest 2.3 times more frequently than the average American.

# Causes of Death

## Underlying and Contributing Factors

When evaluating the cause of death, it is important to examine and understand the underlying and contributing factors that result in the diagnoses of specific diseases and causes of death. While genetics and environmental factors have a part, behavior plays a much larger role in the development of specific diseases that result in death, than was believed in the past.

In 2009-2010, at least 47% of adults in the U.S. had at least one risk factor for heart disease<sup>8</sup>. According to the Centers for Disease Control and Prevention (CDC), there are three prevailing risk factors contributing to the incidence of heart disease including uncontrolled high blood pressure, uncontrolled high low-density lipoproteins cholesterol (LDL) and smoking. (The first two of which can be exacerbated by behavior and the last being solely a behavior.) Similarly, 90% of all lung cancers are the result of smoking tobacco<sup>9</sup> and 80% of all patients diagnosed with COPD are or were smokers<sup>10</sup>.


An examination of the underlying causes of premature death (death before age 75)<sup>7</sup>, for Pendleton County residents shows behavioral factors have a significant effect on the lower life expectancy trends as compared with the nation and state overall. Those behaviors including smoking, alcohol use, and drug/alcohol abuse<sup>11</sup>.

ICD10	Description	Deaths per 100k <sup>7</sup>	Underlying Contributing Factors/100k		
			Tobacco <sup>11</sup>	Alcohol <sup>11</sup>	Drug/Alcohol Abuse <sup>11</sup>
C34.9	Lung Cancer	56.7	51	2	
I21.9	Acute Mia	46.8	9.4		
I25.1   I25.0	Atherosclerotic/Cardiovascular Heart Disease	29.8	6		
J44.9 J43.9	COPD & Emphysema	28.5	22.8		
I50.0	Congestive Heart Failure	21.6	7.1		
V89.2 V49.9 V87.7	Motor Vehicle Accident	20.0		5.6	
X74	Suicide Firearm	17.4			
X42	Accidental Poisoning Narcotics & Psychodysleptics	12.4		6.8	6.8
C18.9	Colon Cancer	11.1		.4	
X44	Accidental Poisoning Unspecified	9.8			5.3
K74.6 K70.3	Liver Cirrhosis	8.2		1.2	
C25.9	Cancer Pancreas	7.9		.3	
C80	Cancer unspecified	7.5			
E14.9	Unspecified Diabetes Mellitus (w/o complications)	7.2			
C50.9	Breast Cancer	6.2		.9	
C71.9	Brain Cancer	5.9		.2	
C61	Prostate Cancer	5.6	1.3	.2	
J18.9	Pneumonia	5.6			
C15.9	Cancer Esophagus	5.6	4.2	4.2	
I11.9	Hypertension	5.2	1		
I64	Stroke	4.9	1		
A41.9	Septicemia, Unspecified	4.6			
I46.9	Heart Attack	4.3	.9		
I26.9	Pulmonary Embolism	4.3			
I42.9	Cardiomyopathy, Unspecified	4.3	.9		
C64	Cancer Kidney	3.9	1.1	.1	
E11.9	Diabetes Mellitus (w/o complication)	3.9			

# Cause of Death


## Years of Potential Life Lost

Years of potential life lost, (YPLL), represent the years of life lost where death occurs before the age of 75, the average annual age of death for Americans. Pendleton County has higher YPLL, as compared with state and top performers in the nation.



For those under the age of 75, the leading cause of death is malignant neoplasms, 148.7 per 100k, followed by ischaemic heart disease at 78.3 per 100k and other diseases of the heart at 44.5 per 100k<sup>7</sup>.

**Cause of Years of Potential Life Lost<sup>7</sup>**  
**Pendleton County 2022**






# Causes of Death

## Cancers


Since the year 1999, cancer of the lungs or bronchus has represented the leading cause of death among persons diagnosed with malignancies in Pendleton County, accounting for 76.9 deaths per 100,000 population<sup>7</sup>.

**Leading Cancer Cause of Death<sup>7</sup>**  
Pendleton County 1999-2020



According to the CDC, smoking can cause cancers almost anywhere in the body including the bladder, blood, cervix, colon, rectum, esophagus, kidney, renal pelvis, liver, lungs, bronchi, trachea, mouth, throat, pancreas, stomach and larynx<sup>13</sup>. Additionally, Lung Cancer has a known association with the inhalation of radon gas. Cancer rates in Pendleton County may be the result of either or both underlying contributing factors. See page 19 for details on tobacco rates. See page 24 for details on radon in Pendleton County.

**Leading Cancer Cause Death<sup>7</sup>**  
Three Rivers District 1999-2020



ICD10	Cancer
C34.9	Lung
C18.9	Colon
C50.9	Breast
C80	Unspecified
C61	Prostate
C25.9	Pancreas
C64	Kidney
C67.9	Bladder
C56	Ovarian
C15.9	Esophagus
C85.9	Lymphoma
C71.9	Brain
C90.0	Multiple Myeloma




# Health Conditions

## Asthma

According to the CDC, 1 in 13 persons in the U.S. has asthma<sup>14</sup>. Most people can control their symptoms and prevent asthma attacks by avoiding triggers and using prescribed medication such as inhalants correctly.

### Asthma Prevalence<sup>14,15</sup>

Pendleton County, Kentucky, U.S. 2016-2018




Pendleton County has a higher-than-average prevalence for asthma at 8.5%, or 1 in 11, when compared with national averages at 7.7%, 1 in 13 people, but is lower than the state averages at 1 in 9 or 11.5% of the population<sup>15</sup>.


***“1 in 11 people in Pendleton County has Asthma”***

Asthma induced ER visits in Pendleton County in 2019 represent an equivalent of 1.7% of the total population or 20% of those with Asthma indicating those with Asthma in

Pendleton Co. may have better control of their condition or have less environmental factors that affect them negatively compared with neighboring communities<sup>16</sup>.

Asthma induced Emergency Room Visits Age Adjusted Fine Particulate Air Matter (Modeled)<sup>16</sup>  
Pendleton County & Kentucky 2008-2019





# Health Conditions


## Chronic Obstructive Pulmonary Disease

Chronic Obstructive Pulmonary Disease, or COPD is a group of pulmonary disease that restricts airflow within the lungs and includes chronic bronchitis and emphysema.

According to the CDC, approximately 18 million Americans have been diagnosed with


COPD and millions more suffer with it but have yet to be diagnosed and receive treatment<sup>17</sup>. People aged 65 and over are more likely to be diagnosed with some form of COPD but it can affect persons of any age.

**COPD induced Hospitalizations Age Adjusted<sup>16</sup>**  
Pendleton County & Kentucky 2008-2019



Nationally, there are 40.4 ER visits per 100 persons and 43.5% of those visits are less than 15 minutes in duration<sup>18</sup>. According to the National Hospital Ambulatory Medical Care Survey: 2018 emergency Department Summary Tables, COPD accounted for 8.1 million ER visits in 2018 or 6.2% of all ER visits nationwide<sup>18</sup>. Pendleton County has seen a two-year average increase of 19% in COPD induced ER visits of which may indicate residents are experiencing more


**COPD induced Emergency Room Visits Age Adjusted<sup>16</sup>**  
Pendleton County & Kentucky 2008-2019




enviornmental triggers, or having issues with access to preventive and acute care.

# Health Conditions


## Heart Attacks



A heart attack or myocardial infarction is a result of reduction or block of blood flow to the heart that results in the damage or death of part of the heart muscle. Coronary Artery Disease (CAD) is the primary cause of myocardial infarction. Coronary Artery Disease is caused by the buildup of plaque and other deposits in the artery(ies) creating narrowing, reducing blood flow and supply to critical body parts/organs.



***“Pendleton County has 1.3 times more heart attacks than KY”***



When overlaying mortality associated with heart attacks in Pendleton County, the percentages of heart attacks resulting in death is high<sup>7</sup>.

*(Note, years with deaths below a specific threshold are suppressed to protect privacy and are not displayed.)*


# Health Conditions

## Diabetes Prevalence

According to the CDC, 37.3 million people have diabetes in the U.S. and 23% are undiagnosed<sup>20</sup>. Diabetes is a chronic life-long disease characterized by high blood sugar resulting in the 7<sup>th</sup> leading cause of death in the U.S. It is the number one cause of kidney failure, lower-limb amputations and adult blindness<sup>21</sup>.

Type 2 diabetes is the most common form of the disease affecting between 90-95% of those diagnosed with diabetes<sup>22</sup>. It may be prevented or delayed with healthy lifestyle changes such as losing weight, eating healthy foods, and being active.

As of 2019, 1 in 13 people in Pendleton County have been diagnosed with diabetes and males have slightly higher rates of diabetes diagnoses than females.



### Risks for Pre-Diabetes and type 2 diabetes<sup>23</sup>

- Overweight
- 45 and older
- Parent or sibling with type 2 diabetes
- Physically active less than 3x/week
- Had gestational diabetes
- African American, Hispanic/Latino American, Pacific Islander, American Indian, or Asian American


# Health Conditions

## Sexually Transmitted Infections

Pendleton County has maintained a significantly lower rate of sexually transmitted infections as the state overall with the incidence of Chlamydia remaining below the state rate consistently with an eight-year trend of 231.7 as compared to the state average at 376.1 per 100k persons<sup>25</sup>.


Likewise, Gonorrhea incidence also remains significantly lower than the state rate averaging 42.2 cases per 100k<sup>25</sup>.

**Rate of sexually transmitted infections<sup>25</sup>**  
Pendleton County & Kentucky 2000-2020



***“47.4% of youths did not use a condom in their last encounter”***

**Sexually Active High School Youths<sup>26</sup>**  
Kentucky 1997-2019



Approximately 40.4% of Kentucky high school students have had at least one sexual encounter<sup>26</sup>. Of those, 7.2% had sexual intercourse with four or more persons and 30.7% are currently sexually active<sup>26</sup>.


# Health Conditions

## Overweight and Obesity – Adults



Adult obesity is defined as a weight that is higher than what is considered healthy for a person’s height. In the years 2021, the adult obesity prevalence was 41.9% of U.S. adults.<sup>27</sup>

Kentucky leads the nation in obesity rates with obesity rates averaging 40-45% of the population overall.




***“67% of Ky counties have lower obesity rates than Pendleton Co.”***

The most recent county level data reflects obesity rates that are higher than the state average prevalence at 40% of the total population in 2020<sup>28</sup> for Pendleton County.

Obesity age-adjusted prevalence (%)


- > 42.8 - 46.2
- > 40.6 - 42.8
- > 38.8 - 40.6
- > 36.6 - 38.8
- 33.2 - 36.6



# Health Conditions


## Overweight and Obesity by Race and Ethnicity

Obesity prevalence affects persons of varying races and ethnicities differently among populations. The prevalence of Self-Reported Obesity, 2019-2021 Behavioral Risk Factor Surveillance System (BRFSS) reflects significant differences in obesity prevalence rates among non-Hispanic, Hispanic, black, white, and Asian populations. Non-Hispanic Asian adults have the lowest obesity prevalence at just 13.8% of the U.S. population as compared with all other races and Hispanic populations<sup>29</sup>.



In Pendleton County, 1 in 3.2 White Adults; 1 in 2.8 Hispanic adults; 1 in 2.5 Black adults are obese and 1 in 7.2 Asian adults are obese<sup>29</sup>.

**Age-adjusted, nationally applied rate of prevalence of adult obesity by race/ethnicity, based on Ky Averages<sup>29</sup>**  
Pendleton County, 2020






# Health Conditions

## Childhood Obesity

Childhood overweight is defined as a Body Mass Index (BMI) at or above the 85<sup>th</sup> percentile and below the 95<sup>th</sup> percentile for children and teens of the same age and sex. Childhood obesity is defined as a BMI at or above the 95<sup>th</sup> percentile for children and teens of the same age and sex. The National Health and Nutrition Examination Survey 2017-March 2020 Prepandemic, indicates an overall obesity prevalence rate among youth aged 2-19 years of 19.7% in the U.S.<sup>30</sup> The survey also found that obesity rates increased among youth as age increased.


**Youth Obesity Rate by Age Group<sup>30</sup>**  
U.S. 2017-March 2020



***“Recent studies show childhood obesity rates are increasing”***

As an attempt to help slow the spread of COVID-19 in Kentucky schools, the last two school years, 2020-2021 and 2021-2022 did not conduct mass health screenings in Kentucky. Therefore, local BMI data for Pendleton County schools is extremely limited and has a very low confidence interval and has not been included in this most recent report. It is worth noting however that the average percent of children considered

**Youth Obesity Rate Trends<sup>31</sup>**  
Kentucky Ages 10-17, 2016-2020



obese in Kentucky schools for ages 10-17 has increased from 19.3% of the population in the school year 2016-2017 to 23.8% in the school year 2019-2020<sup>31</sup>.




# Health Risk Behaviors

## Diet

Healthy behaviors such as a healthy diet and regular physical activity contribute to better short term health outcomes, long term health benefits, and the prevention of chronic diseases such as Type 2 diabetes and heart disease.

Factors affecting healthy behaviors related to diet and exercise include access to and availability of healthy foods and opportunities for exercise. Access is limited in populations that are low income, have limited grocery supermarkets, have a high fast-food concentration and lack of access to transportation. The U.S. Department of Agriculture, Food Access Research Atlas indicators reveals more than half of Pendleton County is categorized as low income<sup>32</sup> which can impact access to healthy foods.



Pendleton County is at a significant disadvantage when compared with residents in areas such as Boone, Kenton, Campbell, and Oldham counties that have little to no low-income areas, and have much higher rates of grocery stores, and full-service restaurants.

### Food establishments<sup>33</sup>

- Groceries – 2
- Full-Service Restaurant – 4
- Fast Food Restaurant – 5
- Convenience Stores – 3
- Famer’s Market – 1


# Health Risk Behaviors

## Exercise

Physical inactivity is a critical factor in the health status of a population. Exercise has been linked to healthy weight and BMI, reduction in various chronic diseases, normal growth and development, better quality of life, and healthy sleep patterns.

***“More than 31% of Pendleton County Residents are inactive”***

The CDC U.S. Diabetes Atlas<sup>34</sup> reports Pendleton County has a self-reported “lack of physical activity” that is statistically lower than the state overall indicating they are less physically active than the average Kentuckian. As of 2020, Pendleton County has a physical inactivity rate of 31.8% of the population reporting no leisure time physical activity in the last 30 days<sup>34</sup>.



The source of data used in the Atlas includes the BRFSS data which defines physical inactivity as those respondents who answered no to the question “During the past month, other than your job, did you participate in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise?”


# Health Risk Behaviors

## Tobacco Use Adults


Smoking remains the single largest leading cause of preventable disease, death, and disability, in the U.S., accounting for more than 480,000 deaths every year, or about 1 in 5 deaths<sup>35</sup>. The Centers for Disease Control and Prevention reports that adult smoking rates have declined in the U.S. from 1 out of nearly every 5<sup>35</sup> adults in 2005 to 1 in 10 as of 2020<sup>36</sup>.

Pendleton County has a significantly higher rate of adults who smoke, with a rate of 25% or 1 in 4 adults<sup>37</sup>.

The financial burden of tobacco use is significant, costing the U.S. billions of dollars each year. Comprehensive tobacco policies have demonstrated positive effects on the rates of tobacco use among



**Adult Smoking Trends<sup>37</sup>**  
Pendleton County, Kentucky, Us 2014-2022<sup>22</sup>



adults and youth alike, recognizing a 10-15% reduction in smoking prevalence where comprehensive tobacco policies have been in place for eight years. (The Journal of Public Health Management and Practice, The Impact of Implementing Tobacco Control Policies: The 2017 Tobacco Control Policy Scorecard).

# Health Risk Behaviors



## Tobacco Use Adolescents

The Centers for Disease Control and Prevention reports that 1 in 4 High School Students used some form of tobacco product in 2022 with the greatest percentage of those using e-cigarettes in the past 30 days at 14.1%<sup>38</sup>.

**“As many as 1 in 6 HS students report using Tobacco”**

Today, girls are more likely to report using of some tobacco product in the last 30 days than boys, a shift since 2018 when more boys reported use than girls, with E-Cigarettes representing 15.4% of tobacco use in girls<sup>38</sup>. The same survey found that similar findings in middle school students reporting use of some form of tobacco product with most of these using e-cigarettes (4.1%)<sup>38</sup>.

Assuming similar applied ratios of tobacco use in the youth population as compared with national averages, middle and high school use could be as high as 1 in 8 or as low as 1 in 14 for middle school students and as high as 1 in 2 or as low as 1 in 4 for high school students in Pendleton County.



# Health Risk Behaviors

## Alcohol Use


In the years 2015-2019 excessive alcohol use was responsible for more than 140,000 deaths in the U.S. each year, or more than 380 deaths per day<sup>39</sup>. In Kentucky men with alcohol related deaths were nearly four times more likely to have chronic alcohol abuse than women<sup>40</sup>.

According to the BRFSS survey results as reported in the 2022 County Health Rankings, Pendleton County has an average number of adults reporting binge or heavy drinking in the last 30 days compared with the state overall<sup>41</sup>.

Excessive or heavy drinking is associated with short and long term health effects including: injuries such as motor vehicle crashes, falls, drownings, and burns; violence including homicide, suicide, sexual assault, and intimate partner violence; alcohol poisoning, risky sexual behaviors including unprotected sex with multiple partners which can result in unwanted pregnancies or sexually transmitted diseases including HIV; and miscarriage or still birth, or fetal alcohol spectrum disorders among pregnant women.

Long term health risks include high blood pressure; heart disease, stroke, liver disease, and digestive problems; cancer of the breast, mouth, throat, esophagus, liver, and colon; learning and memory problems including dementia; mental health problems including depression and anxiety; social problems including lost productivity, family problems and unemployment; alcohol dependence or alcoholism.

**Percent of Adult Population Self-Reporting Excessive Drinking<sup>41</sup>**  
**Pendleton County 2011-2022**



*Excessive alcohol use includes binge drinking, heavy drinking, any alcohol use by persons under 21 years and alcohol used by pregnant women.*

*Binge drinking is defined as 5 or more drinks consumed on a single occasion for men and 4 or more for women, generally within about 2 hours.*

*Heavy drinking is defined as men who consume 15 drinks or more per week and women who consume 8 drinks or more per week.*




# Health Risk Behaviors

## Recreational Drug Use/Abuse

In 2015, the Pendleton County Fiscal Court adopted a comprehensive policy for the creation and operation of a Harm Reduction Syringe Access Exchange Program as permitted by SB192 enacted by the KY General Assembly in 2015. Of the 120 counties in Kentucky, 36 are ranked in the top 100 at risk for serious disease outbreaks including HIV, and 54, or nearly half, are in the top 220 counties across the nation identified at risk<sup>42</sup>. One of the primary objectives of the [Hep C Elimination Plan in Kentucky](#) is to increase and improve equitable access to harm reduction services across social and demographic areas particularly among people who use drugs.


**Kentucky Harm Reduction Syringe Access Programs<sup>43</sup>**



According to the 2021 Kentucky Drug Overdose Deaths Report, the Pendleton County had an age adjusted mortality rate of 92.13 per 100,000 persons<sup>44</sup>.

2021 Drug Overdose Deaths Kentucky<sup>44</sup>


- Suppressed
- 0 per 100k
- 100 - 110 per 100k
- 111 - 125 per 100k
- 126 - 150 per 100k
- Less than 100 per 100k




# HEALTH RISK FACTORS

## Overdose Mortality

In Pendleton County, deaths associated with drug overdose rates increased 51.27% from 55 to 83.2 per 100k, between the 2011-2015 and 2016-2020 time periods<sup>45</sup>. Compared with the Commonwealth and the U.S. Pendleton County mortality from drug overdoses is significantly higher than the Commonwealth and U.S. with the national and Commonwealth rates at 32.5 and 54.1 respectively<sup>45</sup>.



Deaths from opioid specific overdoses also increased significantly, representing a 87.7% increase between the 2011-2015 and 2016-2020 time periods<sup>45</sup>. Concurrently, rates for opioid specific drug overdose deaths in the Commonwealth and U.S. also both increased during the same time period as Pendleton County<sup>45</sup>.






# Environmental Factors

## Air Particulate Matter

Per the CDC, Environmental Public Health Tracking Network, air pollution poses a public health threat affecting potentially millions of people throughout the U.S.<sup>46</sup> Pollution such as dust, dirt, soot, and smoke are one kind of air pollution called fine particulate matter (FPM). Sources of FPM include forest fires, power plants, industries, and automobiles. FPM is associated with compromised health and may lead to breathing problems, worsening of asthma symptoms and/or worsening of some heart conditions, as well as increased low birth weight<sup>47</sup>.

Decreased lung function, chronic bronchitis, asthma, and adverse pulmonary effects are common in populations exposed to high levels of FPM. Long term exposure is associated with premature death of people aged 65 and older even when exposure is at levels below the National Standards<sup>48</sup>. Since 2001 FPM rates in Pendleton County have decreased by 41% to 8.5 PM ug/M<sup>3</sup> in 2018<sup>49</sup>.



Rates below 12 Ug/m<sup>3</sup> are considered acceptable according to the U.S. Environmental Protection


# Environmental Factors

## Radon

Radon is a gas formed naturally in the environment with the breakdown of radioactive metals. People are always exposed to radon because it comes from the earth, but some areas have higher concentrations of the gas than others. The gas can enter your home

through cracks and gaps.

Kentucky has an average radon level of 5.4 picocuries per liter of air (pCi/L)<sup>50</sup>. The



Environmental Protection Agency indicates remediation should be taken for levels exceeding 4pCi/L in a home<sup>51</sup>. Higher radon levels are associated with increased risk of lung cancer and is the second leading cause of all lung cancers in the U.S.


***“Pendleton Co. average radon levels for homes tested is 3.1pCi/L”***

Of the homes tested, the average indoor radon level is 3.1 pCi/L as compared to 1.3pCi/L nationally<sup>52</sup>. Pendleton County homes have higher indoor radon levels than

the nation. Of the homes measured in Pendleton County, 19% have radon levels of 4pCi/L or higher<sup>52</sup>.

### Indoor Radon Levels measured in pCi/L<sup>52</sup>

#### Pendleton County, 2022



# End Notes

## References and Data Sources

1. Institute of Health Metrics Evaluation, US Health Map, Viz Hub, Life Expectancy, 2000-2019 as accessed at <https://vizhub.healthdata.org/subnational/usa>
2. Institute of Health Metrics Evaluation, 1980-2014 United States Life Expectancy and Age-specific Mortality Risk by County 1980-2014. Seattle, United States of America, institutes for Health Metrics and Evaluation (IHME), 2017, as accessed at <https://ghdx.healthdata.org/record/ihme-data/united-states-life-expectancy-and-age-specific-mortality-risk-county-1980-2014>
3. U.S. Census Bureau, American Fact Finder, 2020 ACS 5 Year Estimates, Table DP03 – Selected Economic Characteristics
4. U.S. Census Bureau, American Fact finder, 2020 ACS 5 Year Estimates, Table S1501 – Educational Attainment
5. U.S. Census Bureau, American Fact Finder, 2020 ACS 5 Year Estimates, Table S2404 – Industry by Sex for the Full-Time, Year-Round Civilian Employed Population 16 Years and Over
6. Centers for Disease Control and Prevention, National Center for Health Statistics, Leading Causes of Death, 1900-1998 as accessed at [https://www.cdc.gov/nchs/data/dvs/lead1900\\_98.pdf](https://www.cdc.gov/nchs/data/dvs/lead1900_98.pdf)
7. Centers for Disease Control and Prevention, National Center for Health Statistics. National Vital Statistics System, Mortality 1999-2020 on CDC WONDER Online Database, released in 2021. Data are from the Multiple Cause of Death Files, 1999-2020, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program. Accessed at <http://wonder.cdc.gov/ucd-icd10.html> on Dec 5, 2022, 2:24:37 PM
8. Fryar CD, Chen T, Li X. Prevalence of uncontrolled risk factors for cardiovascular disease: United States, 1999–2010. NCHS data brief, no 103. Hyattsville, MD: National Center for Health Statistics. 2012. As accessed at <https://www.cdc.gov/nchs/data/databriefs/db103.htm>
9. U.S. Department of Health and Human Services. The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General. Atlanta: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2014 As accessed at: [https://www.cdc.gov/tobacco/data\\_statistics/sgr/50th-anniversary/index.htm#report](https://www.cdc.gov/tobacco/data_statistics/sgr/50th-anniversary/index.htm#report)
10. The COPD Foundation, Understanding COPD, What Causes COPD. As accessed at <https://www.copdfoundation.org/What-is-COPD/Understanding-COPD/What-Causes-COPD.aspx>
11. See Appendix A
12. County Health Rankings, Pendleton County, Kentucky, Premature Death, 2022.
13. The Centers for Disease Control and Prevention, Smoking and Cancer, May 2022, as accessed at: <https://www.cdc.gov/tobacco/campaign/tips/diseases/cancer.html>
14. The Centers for Disease Control and Prevention, National Current Asthma Prevalence, 2020, as accessed at: [https://www.cdc.gov/asthma/most\\_recent\\_national\\_asthma\\_data.htm](https://www.cdc.gov/asthma/most_recent_national_asthma_data.htm)
15. The Centers for Disease Control and Prevention, State Maps for Asthma Prevalence by Six-Level Urban-Rural Classification, 2016-2018, as accessed at: <https://www.cdc.gov/asthma/national-surveillance-data/asthma-prevalence-state-classification.htm>
16. Kentucky Environmental Health Tracking, Asthma induced Emergency Room Visits Age Adjusted and Fine Particulate Air Matter Modeled as access at: <https://kyibis.mc.uky.edu/ehl/>
17. Centers for Disease Control and Prevention, National Trends COPD, As accessed at: <https://www.cdc.gov/copd/data-and-statistics/national-trends.html>
18. National Hospital Ambulatory Medical Care Survey: 2018 Emergency Department Summary Tables as accessed at: [https://www.cdc.gov/nchs/data/nhamcs/web\\_tables/2018-ed-web-tables-508.pdf](https://www.cdc.gov/nchs/data/nhamcs/web_tables/2018-ed-web-tables-508.pdf)

19. Kentucky Environmental Health Tracking, Number of Hospitalizations for Heart Attack by County, as access at: <https://kyibis.mc.uky.edu/ehl/>
20. Centers for Disease Control and Prevention, Fast Facts on Diabetes as accessed at: <https://www.cdc.gov/diabetes/data/statistics-report/index.html>
21. Centers for Disease Control and Prevention, Diabetes Basics, What is Diabetes, as accessed at: <https://www.cdc.gov/diabetes/basics/diabetes.html>
22. Center for Disease Control and Prevention, Diabetes Basics, Type 2 Diabetes, as accessed at: <https://www.cdc.gov/diabetes/basics/type2.html>
23. Centers for Disease Control and Prevention, About Prediabetes & Type 2 Diabetes as accessed at: <https://www.cdc.gov/diabetes/prevention/about-prediabetes.html>
24. Centers for Disease Control and Prevention, United States Diabetes Surveillance System, Surveillance, Diabetes Prevalence, Pendleton County as accessed at: <https://gis.cdc.gov/grasp/diabetes/DiabetesAtlas.html>
25. Centers for Disease Control and Prevention, Atlas Plus, HIV, Hepatitis, STD, TB, SDOH, 2020 as accessed at: <https://gis.cdc.gov/grasp/nchhstpatlas/tables.html>
26. Youth Behavior Risk Surveillance Survey 2019, Kentucky Results, as accessed at: <https://nccd.cdc.gov/youthonline/App/Results.aspx?LID=KY>
27. Centers for Disease Control and Prevention, Adult Obesity, as accessed at: <https://www.cdc.gov/obesity/data/adult.html>
28. Centers for Disease Control and Prevention, Explore Places Interactive Map, Obesity Rates 2020, as accessed at: [https://experience.arcgis.com/experience/dc15b033b88e423d85808ce04bd7a497/page/Health-Outcomes/?data\\_id=dataSource\\_17-PLACES\\_LocalData\\_for\\_BetterHealth\\_5583%3A1860&views=Obesity](https://experience.arcgis.com/experience/dc15b033b88e423d85808ce04bd7a497/page/Health-Outcomes/?data_id=dataSource_17-PLACES_LocalData_for_BetterHealth_5583%3A1860&views=Obesity)
29. Centers for Disease Control and Prevention, Overweight & Obesity, Data & Statistics, Adult Obesity Prevalence Maps as accessed at: <https://www.cdc.gov/obesity/data/prevalence-maps.html#race>
30. National Health and Nutrition Examination Survey 2017-March 2020 Prepandemic Data Files Development of Files and Prevalence Estimates for Selected Health Outcomes, Youth Obesity, As accessed at: <https://stacks.cdc.gov/view/cdc/106273>
31. Robert Wood Johnson Foundation, State of Childhood Obesity, Explore State Data, Kentucky, Children ages 10-17, as accessed at: <https://stateofchildhoodobesity.org/state-data/?state=ky>
32. U.S. Economic Research Service, U.S. Department of Agriculture, Food Access Research Atlas as accessed at: <https://www.ers.usda.gov/data-products/food-access-research-atlas/go-to-the-atlas/>
33. U.S. Department of Agriculture, Food Environment Atlas, Pendleton County. As accessed at: <https://www.ers.usda.gov/data-products/food-environment-atlas/go-to-the-atlas/>
34. Centers for Disease Control and Prevention, Explore Places Interactive Map, Physical Inactivity Rates 2020, as accessed at: [https://experience.arcgis.com/experience/dc15b033b88e423d85808ce04bd7a497/page/Health-Outcomes/?data\\_id=dataSource\\_17-PLACES\\_LocalData\\_for\\_BetterHealth\\_5583%3A1140&views=Obesity](https://experience.arcgis.com/experience/dc15b033b88e423d85808ce04bd7a497/page/Health-Outcomes/?data_id=dataSource_17-PLACES_LocalData_for_BetterHealth_5583%3A1140&views=Obesity)
35. U.S. Department of Health and Human Services. The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General. Atlanta: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2014. As accessed at: [https://www.ncbi.nlm.nih.gov/books/NBK179276/pdf/Bookshelf\\_NBK179276.pdf](https://www.ncbi.nlm.nih.gov/books/NBK179276/pdf/Bookshelf_NBK179276.pdf)
36. Centers for Disease Control and Prevention, Smoking & Tobacco Use, Data and Statistics, Fast Facts and Fact Sheets, Smoking and Cigarettes as accessed at: [https://www.cdc.gov/tobacco/data\\_statistics/fact\\_sheets/fast\\_facts/index.htm?CDC\\_AA\\_refVal=https%3A%2F%2Fwww.cdc.gov%2Ftobacco%2Fdata\\_statistics%2Ffact\\_sheets%2Findex.htm](https://www.cdc.gov/tobacco/data_statistics/fact_sheets/fast_facts/index.htm?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Ftobacco%2Fdata_statistics%2Ffact_sheets%2Findex.htm)
37. County Health Rankings, Kentucky, Adult Smoking, 2022 as accessed at: <https://www.countyhealthrankings.org/explore-health-rankings/kentucky/pendleton?year=2022>
- 38.

39. Centers for Disease Control and Prevention, Tobacco, Data and Statistics, Fact Sheets, Youth and Tobacco Use, as accessed at: [https://www.cdc.gov/tobacco/data\\_statistics/fact\\_sheets/youth\\_data/tobacco\\_use/index.htm](https://www.cdc.gov/tobacco/data_statistics/fact_sheets/youth_data/tobacco_use/index.htm)
40. Centers for Disease Control and Prevention, Alcohol and Public Health, Data on Excessive Drinking as accessed at: <https://www.cdc.gov/alcohol/data-stats.htm>
41. Centers for Disease Control and Prevention, Alcohol-Related Disease Impact (ARDI), Alcohol-attributable deaths, 2015-2019 as accessed at: [https://nccd.cdc.gov/DPH\\_ARDI/Default/Default.aspx](https://nccd.cdc.gov/DPH_ARDI/Default/Default.aspx)
42. County Health Rankings, Pendleton County, Kentucky, Excessive Drinking, 2022 as accessed at: <https://www.countyhealthrankings.org/explore-health-rankings/kentucky/pendleton?year=2022>
43. Centers for Disease Control and Prevention, Vulnerable Counties and Jurisdictions Experiencing or At-Risk of Outbreaks, as accessed at: <https://www.cdc.gov/pwid/vulnerable-counties-data.html>
44. Kentucky Cabinet for Health and Family Services, Department for Public Health, Division of Epidemiology and health Planning, HIV/AIDS Branch, Syringe Exchange Programs as accessed at: <https://www.chfs.ky.gov/agencies/dph/dehp/hab/Pages/kyseps.aspx>
45. Recovery Ecosystem Index as accessed at: [https://rsconnect.norc.org/recovery\\_ecosystem\\_index/](https://rsconnect.norc.org/recovery_ecosystem_index/)
46. Kentucky Office of Drug Control Policy, 2021 Overdose Fatality Report as accessed at: [https://odcp.ky.gov/Reports/2021%20Overdose%20Fatality%20Report%20\(final\).pdf](https://odcp.ky.gov/Reports/2021%20Overdose%20Fatality%20Report%20(final).pdf)
47. Centers for Disease Control and Prevention, National Environmental Public Health Tracking, Explore Environmental Health Data, Air Quality as accessed at: <https://www.cdc.gov/nceh/tracking/topics/AirQuality.htm>
48. Particle Pollution, Fine particulate Air Matter, Centers for Disease Control and Prevention. As accessed at: [https://www.cdc.gov/air/particulate\\_matter.html](https://www.cdc.gov/air/particulate_matter.html)
49. The Harvard T.H. Chan School of Public Health, Harvard Data Science Initiative, Press Release, 2017. As accessed at: <https://www.hsph.harvard.edu/news/press-releases/u-s-seniors-air-pollution-premature-death/>
50. Centers for Disease Control and Prevention National Environmental Public Health Tracking Network, Fine Particulate Air Matter as accessed at: <https://ephtracking.cdc.gov/DataExplorer/>
51. American Lung Association, Radon Testing Disparities, Kentucky, Radon Testing in the State of Kentucky as accessed at: <https://www.lung.org/getmedia/5e0f8076-92c9-482e-b658-588d97b0b7e5/Radon-Testing-Disparity-report-for-Kentucky-2022-08-14>
52. U.S. Environmental Protection Agency, A Citizen's Guide to Radon, Protecting yourself and your Family from Radon, as accessed at: [https://www.epa.gov/sites/default/files/2016-12/documents/2016\\_a\\_citizens\\_guide\\_to\\_radon.pdf](https://www.epa.gov/sites/default/files/2016-12/documents/2016_a_citizens_guide_to_radon.pdf)
53. University of Kentucky, Kentucky Geological Survey, Radon as accessed at: <https://www.uky.edu/KGS/radon/>

## Appendix A

ICD10	Tobacco Association	Alcohol Association
C34.9	<a href="https://pubmed.ncbi.nlm.nih.gov/16100660/">https://pubmed.ncbi.nlm.nih.gov/16100660/</a>	<a href="https://www.livescience.com/27259-alcohol-causes-cancer-deaths.html">https://www.livescience.com/27259-alcohol-causes-cancer-deaths.html</a>
I21.9	<a href="https://www.ncbi.nlm.nih.gov/books/NBK53012/">https://www.ncbi.nlm.nih.gov/books/NBK53012/</a>	
J44.9	<a href="https://www.cdc.gov/tobacco/data_statistics/fact_sheets/health_effects/effects_cig_smoking/index.htm">https://www.cdc.gov/tobacco/data_statistics/fact_sheets/health_effects/effects_cig_smoking/index.htm</a>	
I25.1	<a href="https://www.ncbi.nlm.nih.gov/books/NBK53012/">https://www.ncbi.nlm.nih.gov/books/NBK53012/</a>	
I50.0	<a href="https://www.ncbi.nlm.nih.gov/books/NBK53012/">https://www.ncbi.nlm.nih.gov/books/NBK53012/</a>	
V89.2		<a href="https://www.cdc.gov/motorvehiclesafety/impaired_driving/impaired_drv_factsheet.html#statistics">https://www.cdc.gov/motorvehiclesafety/impaired_driving/impaired_drv_factsheet.html#statistics</a>
C50.9		<a href="https://www.livescience.com/27259-alcohol-causes-cancer-deaths.html">https://www.livescience.com/27259-alcohol-causes-cancer-deaths.html</a>
X44		CDC wonder calculated by # of deaths caused by drug and alcohol induced from total poisoning deaths
A41.9		
C18.9		<a href="https://www.livescience.com/27259-alcohol-causes-cancer-deaths.html">https://www.livescience.com/27259-alcohol-causes-cancer-deaths.html</a>
J18.9		
X70/X74		
X42		CDC wonder calculated by # of deaths caused by drug and alcohol from total poisoning deaths
E14.9		
J44.0	<a href="https://www.cdc.gov/tobacco/data_statistics/fact_sheets/health_effects/effects_cig_smoking/index.htm">https://www.cdc.gov/tobacco/data_statistics/fact_sheets/health_effects/effects_cig_smoking/index.htm</a>	
K70.3		
C80		
J43.9	<a href="https://www.cdc.gov/tobacco/data_statistics/fact_sheets/health_effects/effects_cig_smoking/index.htm">https://www.cdc.gov/tobacco/data_statistics/fact_sheets/health_effects/effects_cig_smoking/index.htm</a>	
I64	<a href="https://www.ncbi.nlm.nih.gov/books/NBK53012/">https://www.ncbi.nlm.nih.gov/books/NBK53012/</a>	
I11.9	<a href="https://www.ncbi.nlm.nih.gov/books/NBK53012/">https://www.ncbi.nlm.nih.gov/books/NBK53012/</a>	
C25.9		<a href="https://www.livescience.com/27259-alcohol-causes-cancer-deaths.html">https://www.livescience.com/27259-alcohol-causes-cancer-deaths.html</a>
C92		
C64	<a href="https://www.cancer.net/cancer-types/kidney-cancer/risk-factors-and-prevention#:~:text=Smoking.,Gender.">https://www.cancer.net/cancer-types/kidney-cancer/risk-factors-and-prevention#:~:text=Smoking.,Gender.</a>	
I46.9		